



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$x^{p^n} \equiv x \pmod{p^n}$  has  $p$  and  $p$  solutions  $\pmod{p^n}$ . Hence the congruence defines the Galois field of order  $p^n$  if and only if  $n=1$ .

---

#### AVERAGE AND PROBABILITY.

---

148. Proposed by M. C. RORTY, Boston, Mass.

Assuming  $n$  points to fall at random upon a circle of circumference  $a$ , what is the probability of  $m$  or more points falling within a length  $b$  upon this circumference.

NOTE. This problem has practical application in determining the probability of accidental rushes of telephone calls as distinct from those rushes which are due to commercial causes. The solution for  $m$  or more points falling within a specified length  $b$  is known. The problem presented above differs from this in that a solution is required for *any* length  $b$ .

149 Proposed by L. C. WALKER, A.M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

Three points are taken at random on the convex surface of a right cone. Find the probability that the section of the cone made by the plane passing through them is a complete ellipse.

---

#### MISCELLANEOUS.

---

142. Proposed by E. A. WELLS, Franklin College, New Athens, O.

Find a general expression for the value of  $\theta$  such that when  $\theta$  is one of the acute angles of a right triangle, the three sides of the triangle will be commensurable.

---

#### NOTES.

---

Dr. C. N. Haskins has been appointed Instructor in Mathematics at Yale University.

Dr. A. B. Pierce has been appointed Instructor in Mathematics at the University of Michigan.

Dr. G. B. Halsted has been elected Professor of Mathematics at Kenyon College, Gambier, O.

Mr. A. C. Minear has been appointed Professor of Mathematics in the University of Southern California, to succeed Mr. Paul Arnold, resigned.

Dr. H. C. DeMott, Principal of the Preparatory Department of the Illinois Wesleyan University, has been promoted to the University chair of mathematics.

In line 14 of Professor Himel's Note in the October number, read "Thus, in the above example, if 2 be true."